

Acetaminophen (Tylenol) Dosage Table (for Fever and Discomfort/Pain)

• Call the office immediately if your child is under 2 months old or has not received 2-month vaccines and has a rectal temperature of 100.4F (38C) or above.

Weight in pounds (kg) →	6-11 (2.7-5.3)	12-17 (5.4-7.9)	18-23 (8-10.6)	24-35 (10.7-15.9)	36-47 (16-21.2)	48-59 (21.3-26.6)	60-71 (26.7-31.9)	72-95 (32-42.6)	96+ (42.7+)
Liquid Suspension: 160mg/5mL	1.25 mL (1/4 tsp)	2.5 mL (1/2 tsp)	3.75 mL (3/4 tsp)	5 mL (1 tsp)	7.5 mL (1 ½ tsp)	10 mL (2 tsp)	12.5 mL (2 ½ tsp)	15 mL (3 tsp)	20 mL (4 tsp)
Chewable <u>80mg</u> Tablet	--	--	1 ½ tabs	2 tabs	3 tabs	4 tabs	5 tabs	6 tabs	8 tabs
Chewable <u>160mg</u> Tablet	--	--	--	1 tab	1 ½ tabs	2 tabs	2 ½ tabs	3 tabs	4 tabs
Adult 325mg Tablet	--	--	--	--	--	1 tab	1 tab	1 ½ tabs	2 tabs
Adult 500mg Tablet	--	--	--	--	--	--	--	1 tab	1 tab

- **MEASURING the DOSAGE:** Syringes and droppers are more accurate than teaspoons. If possible, use the syringe or dropper that comes with the medicine. If not, medicine syringes are available at pharmacies. If you use a teaspoon, it should be a measuring spoon. Remember that 1 level teaspoon equals 5 ml, and that 1/2 teaspoon equals 2.5 ml.
- **FREQUENCY:** Repeat every 4-6 hours as needed. Do not give more than 5 times a day.

Ibuprofen (Motrin, Advil) Dosage Table (for Fever and Discomfort/Pain)

• Do NOT give to children under 6 months of age

Weight in pounds (kg) →	12-17 (5-7.4)	18-21 (7.5-9.9)	22-32 (10-14.9)	33-43 (15-19.9)	44-54 (20-24.9)	55-65 (25-29.9)	66-88 (30-39.9)	88+ (40+)
Infant Drops: 50mg/1.25mL	1.25 mL (1/4 tsp)	1.875 mL	2.5 mL (1/2 tsp)	3.75 mL (3/4 tsp)	5 mL (1 tsp)	--	--	--
Children's Suspension: 100mg/5mL	2.5 mL (1/2 tsp)	3.75 mL (3/4 tsp)	5 mL (1 tsp)	7.5 mL (1 ½ tsp)	10 mL (2 tsp)	12.5 mL (2 ½ tsp)	15 mL (3 tsp)	20 mL (4 tsp)
Chewable <u>50mg</u> Tablet	--	--	2 tabs	3 tabs	4 tabs	5 tabs	6 tabs	8 tabs
Chewable <u>100mg</u> Tablet	--	--	1 tab	1 ½ tabs	2 tabs	2 ½ tabs	3 tabs	4 tabs
Adult 200mg Tablet	--	--	--	--	1 tab	1 tab	1 ½ tabs	2 tabs

- **MEASURING the DOSAGE:** Syringes and droppers are more accurate than teaspoons. If possible, use the syringe or dropper that comes with the medicine. If not, medicine syringes are available at pharmacies. If you use a teaspoon, it should be a measuring spoon. Remember that 1 level teaspoon equals 5 ml, and that 1/2 teaspoon equals 2.5 ml.
- **FREQUENCY:** Repeat every 6-8 hours as needed.